



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Reconsideration of interferon treatment for viral diseases: Lessons from SARS, MERS, and COVID-19

Dan Ma^{a,b,1}, Ximin Wang^{c,1}, Min Li^c, Chujiào Hu^{b,*}, Lei Tang^{b,*}

^a Department of Hematology, Affiliated Hospital of Guizhou Medical University, Guiyang 550004, GuiZhou, China

^b Guizhou Provincial Engineering Technology Research Center for Chemical Drug R&D, Guizhou Medical University, Guiyang 550004, GuiZhou, China

^c Department of Pharmacy, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, Hubei, China

ARTICLE INFO

Keywords:

Interferon
Coronavirus
Pneumonia
Viral replication
Efficacy

ABSTRACT

Periodic pandemics of coronavirus (CoV)-related pneumonia have been a major challenging issue since the outbreak of severe acute respiratory syndrome (SARS) in 2002 and Middle East respiratory syndrome (MERS) in 2012. The ongoing pandemic of CoV disease (COVID-19) poses a substantial threat to public health. As for the treatment options, only limited antiviral agents have been approved hitherto, and clinicians mainly focus on currently available drugs including the conventional antiviral interferons (IFNs). In clinical practice, IFNs, when used either alone or in combination with ribavirin and/or lopinavir/ritonavir, have shown promising outcomes, to some extent, in SARS-CoV or MERS-CoV treatment. Although the efficacy and safety of IFNs in COVID-19 treatment remain unclear, their possible use merits further evaluation. We present a review that summarizes current evidence of IFN treatment for COVID-19 and elaborates on other challenges in terms of the timing of IFN treatment initiation, treatment duration, and IFN type to be used. The review findings suggested that IFN acts by directly inhibiting viral replication and activating immune cell subsets. However, there is a lack of well-designed and controlled clinical trials providing firm evidence for the efficacy or safety of IFN therapy for CoVs. Additionally, critically ill patients with multiple immunosuppression-associated comorbidities may not benefit from IFN therapy, necessitating screening of those patients who would most benefit from IFN treatment.

1. Introduction

The ongoing pandemic of coronavirus (CoV) disease (COVID-19), which started in December 2019, was caused by an outbreak of infection due to novel CoV, which was later officially named by the World Health Organization (WHO) as severe acute respiratory syndrome CoV-2 (SARS-CoV-2) [1]. Early reports of the highly pathogenic COVID-19 included the SARS epidemic in late 2002 and Middle East respiratory syndrome (MERS) in 2012 [2,3]. Owing to their unprecedented speed and range of transmission, CoV infections are associated with a high fatality rate and are widely distributed across geographic regions, both of which pose a major threat to healthcare systems and economies.

One of the major challenging issues in the management of COVID-19 is that effective therapeutic regimens are not yet fully established. Even though new drugs such as PaxlovidTM (nirmatrelvir/ritonavir) are approved for emergency use, data on their actual effect are insufficient and controversial [4]. Most of the currently available treatment options

for COVID-19, such as interferons (IFNs) and other antiviral agents, are extrapolated from evidence accumulated during the SARS and MERS epidemics [5]. Physiologically, IFNs are important components of our innate and adaptive immune systems, and exogenously administered IFNs are effective in reducing viral replication and disease severity [6]. Therefore, IFNs are one of the most widely used therapeutic regimens for COVID-19.

This review provides an overview of the theoretical basis of IFN therapy for COVID-19 as well as evidence from in vitro, animal, and clinical studies (Fig. 1). By compiling accumulated information, our review also elaborates on the factors that could significantly affect therapeutic outcomes, such as the route and approach of IFN administration, the timing of IFN initiation, and the duration of IFN treatment, and so on. Given the scenario of the ongoing aggressive COVID-19 pandemic, the findings of this review may help pave the way for more opportunities and consideration of IFN treatment for CoV infections, especially in future clinical trials.

* Corresponding authors at: Guizhou Medical University, Guiyang 550004, China.

E-mail addresses: huchujiao123@163.com (C. Hu), tanglei19742022@163.com (L. Tang).

¹ Contributed equally to this work.

2. COVID-19

CoVs are a group of enveloped, positive-sense single-stranded RNA viruses that are named after their corona-like morphological appearance [7]. They are zoonotic pathogens and can be transmitted from animals to humans, wherein they cause a spectrum of illnesses ranging from being asymptomatic to severe and often fatal respiratory and enteric diseases in their new hosts. In the past decades, pneumonia caused by the highly pathogenic CoV emerged periodically in different areas worldwide and posed substantial challenges to public health. Since the emergence of SARS, the WHO has recorded 8000 or more cases of SARS-CoV infection and approximately 800 associated deaths [8,9]. As of August 2022, the WHO has recorded 2591 laboratory-confirmed MERS cases in 27 countries, with 894 associated deaths [10]. The cumulative number of COVID-19 cases in 196 countries and regions as of October 07, 2022, was 620,471,620, and the total number of associated deaths was 6,554,523 [11].

A patient with COVID-19 may remain asymptomatic in the early stage of the disease until symptoms such as severe pneumonia, dyspnea, septic shock, multiorgan insufficiency and even death occur [12,13]. Initially, most of the infected patients experience flu-like symptoms, commonly including fever, chills, headaches and cough. Some of the cases then rapidly progress into severe pneumonia and acute respiratory distress, and many people with such conditions may require mechanical ventilation [14]. The mortality rate is higher among elderly and comorbid patients, which may be attributed to their compromised immune system and the inhibition of type I IFN signaling by their antibodies, resulting in a significantly increased viral load [15–18]. Hence, COVID-19 is a severe health problem that requires great attention from the public and clinical investigators.

3. Theoretical basis of IFN treatment for COVID-19

IFNs are classified into type I IFNs ($-\alpha$, $-\beta$, $-\delta$, $-\epsilon$, $-\zeta$, $-\kappa$, $-\tau$, and $-\omega$), type

II IFN ($-\gamma$), and type III IFNs ($-\lambda 1$, $-\lambda 2$, $-\lambda 3$, and $-\lambda 4$) according to the type of receptors through which they initiate signaling [19]. Traditionally, IFNs, especially type I IFNs, have been widely used to treat viral hepatitis, malignancies such as leukemia and renal cell carcinoma, and multiple sclerosis [20]. Their efficacy against CoVs is being tested in an ongoing clinical trial, based on their general ability to inhibit viral replication and activate immune cell populations to eliminate viral infections.

IFNs predominantly participate in the host defense function against viral invasion. CoVs first fuse with the cellular membrane and then interact with innate pattern recognition receptors to enter the cells. After viral invasion, considerable levels of type I IFNs and moderate levels of type III IFNs are rapidly produced by stromal cells or professional antigen-presenting cells, triggering strong antiviral innate immune responses [12,21–23]. IFNs can upregulate the expression of more than 300 IFN-stimulated genes (ISGs), leading to the secretion of potent antiviral proteins [24], thus providing first-line defense against diverse pathogens (Fig. 2). Molecules of the type I IFN pathway play critical roles in protecting developing fetuses and resisting SARS-CoV-2 infection in the placenta [25]. Indeed, dysregulation of the IFN response can increase viral virulence [19]. In MERS-CoV-infected mice, blockade of IFN-I signaling delayed clearance of the virus, increased neutrophil infiltration, and impaired MERS-CoV-specific T-cell responses [26]. In patients with moderate COVID-19, physiological levels of IFN- α exerted immunoregulatory effects by suppressing SARS-CoV-2 replication in human airway cells [27]. Reportedly, in patients with COVID-19, SARS-CoV-2 can block the production of IFNs and reduce effective adaptive immune responses by activating the innate immune system through the renin-angiotensin and kinin-kallikrein pathways [28]. Consistently, one patient who had a surging production of IFN- α had a successful clearance of MERS-CoV, whereas another patient who had a decreased production of IFN- α died, which confirms the therapeutic role of IFNs [29]. Type I IFNs not only act as direct antiviral messengers but also promote phagocytosis of macrophages and apoptosis of natural killer cells [19].




Fig. 1. Current research prospects of interferon treatment for COVID-19.




Fig. 2. Antiviral mechanism of interferon. Abbreviations: IFNs, interferons; ISGs, IFN-stimulated genes; PRR, pattern recognition receptor; APCs, antigen-presenting cells.

IFNs are ideal candidates as broad-spectrum antivirals because they exert direct and extensive effects on viral replication together with critical effects on the development and activation of immune cell subsets [19].

CoVs, however, encode an unusually large number of proteins and other factors that facilitate their survival and replication in the host despite the presence of effective host defense mechanisms, making the virus highly pathogenic [30]. In the past decades, tremendous progress has been achieved toward the understanding of anti-IFN activities against CoVs in humans [30,31]. Like most metazoan viruses, CoVs can attenuate the antiviral IFN signals through a multitude of passive and active mechanisms [32]. The nucleocapsid protein of SARS-CoV inhibits IFN- β production and counteracts host defense systems [33]. MERS-CoV can inhibit the body's ability to recognize the virus, antagonize induction of IFN responses, and block ISGs production until peak viral titers are achieved [6,34,35]. Consequently, antigen presentation of the virus and adaptive Th-1 immune responses against the virus diminish with a depressed IFN response [36]. Thus, CoVs undergo immune evasion in the host probably by reducing the IFN response.

Fortunately, the antagonistic activity of the virus against the IFN response, which is imperfect, usually delays the response induction rather than completely suppressing it [32]. Additionally, SARS-CoV and MERS-CoV are sensitive to exogenously administered IFNs [21,37]. In a genome-wide association study, analysis of the viral phenotype in 81 patients with COVID-19 revealed associations with IFN pathways, including SARS-CoV and innate immunity [38]. Notably, MERS-CoV is substantially more sensitive to IFNs than SARS-CoV as demonstrated in a cell culture study [39]. Based on the aforementioned rational and reliable evidence, recombinant IFNs are potentially suitable candidates for COVID-19 treatment.

4. Nonhuman studies

IFNs have been used for COVID-19 treatment based on evidence from (i) *in vitro* experiments; (ii) animal models; and (iii) limited clinical data

for SARS, MERS, or COVID-19. As mentioned above, IFN response plays a key role in host defense mechanisms against viral invasion. Evidence from *in vitro* and animal models has confirmed that IFNs improve treatment outcomes in CoV infection.

4.1. *In vitro* study

The cytopathic effects (CPEs) of SARS-CoV can be inhibited by IFN- α , IFN- β , IFN- γ , or IFN- λ *in vitro* [40–45]. Of these, IFN- β has the highest potency when compared with IFN- γ and IFN- α [36]. Cinatl et al. assessed the inhibitory potential of recombinant IFNs against two clinical isolates of SARS-CoV in Vero and Caco2 cells and reported that IFN- β had a higher anti-SARS-CoV activity than the other IFN types [40], consistent with the findings of several studies [42,46–48]. This finding confirms the prophylactic protective and antiviral effects of IFN- β in SARS-CoV infection. Furthermore, substances that stimulate IFN production in human peripheral blood mononuclear cells, such as CpG oligodeoxynucleotide (BW001), could produce highly protective effects against SARS-CoV infection [49]. Some studies have indicated that IFN- α - β combined with ribavirin [44,50] and IFN- β combined with IFN- γ can synergistically exert inhibitory effects on the replication potential of SARS-CoV [43,51]. For instance, IFN- β combined with IFN- γ inhibited SARS-CoV plaque formation by 30-fold and SARS-CoV replication by 3000-fold compared with the data obtained when these drugs were used as monotherapy [43,51]. Additionally, IFN- γ (100 IU/ml) combined with IFN- β 1b (at a relatively low concentration [137 IU/ml]) adequately restricted viral replication by 93%, whereas monotherapy of these drugs could only marginally abate viral replication [51].

IFNs (α , β , γ , and λ) are considered effective against MERS-CoV *in vitro* [36]. Among them, IFN- β has the strongest biological activity against MERS, reducing viral titers very effectively with an IC_{50} value of 39 IU/ml [52]. IFN- α , similar to pegylated (PEG)-IFN- α , at a dose of 30 ng/ml could completely inhibit CPEs and reduce viral RNA levels in Vero cells [53]. Recently, the protective effect of IFN- λ gained recognition. Jeon et al. quantified the mRNA levels after MERS-CoV infection

and reported that IFN- λ is predominantly produced in the respiratory epithelium, and IFN- λ 4, which has the highest potency to induce a response among all IFN types, demonstrated a remarkable suppressive effect on MERS-CoV replication and ISG expression [54]. Moreover, the type of drug combination was also considered for improved management of patients with MERS. Falzarano et al. reported that IFN- α 2b combined with ribavirin, when administered at a lower concentration, could decrease the replication potential of MERS-CoV compared with monotherapy of the drugs when administered at a higher concentration individually; this finding indicates that the combination regimen facilitated a reduction in the drug concentration by 8-fold and 16-fold, respectively, to obtain a comparable endpoint with monotherapy [55]. Intriguingly, immunosuppressants combined with IFNs were highly effective against the virus. Cyclosporine combined with IFN- α 1 reduced the viral titer more saliently than monotherapy of the drugs [56]. Additionally, mycophenolic acid, when combined with IFN- β 1b, decreased the EC₅₀ of IFN- β 1b [57]. Nevertheless, contradictory results have also been reported. Sheahan et al. reported that lopinavir/ritonavir (LPV/RTV) did not boost the antiviral ability of IFN- β and had no effect on viral replication or severe lung pathology, although the combination regimen (LPV + RTV + IFN- β) could improve pulmonary function [48].

Treatment with IFN- α or IFN- β at a dose of 50 IU/ml reduced SARS-CoV-2 titers by 3.4 log or more than 4 log, respectively, in Vero cells [58]. When compared with SARS-CoV, SARS-CoV-2 has greater sensitivity to IFN- α [59]. Furthermore, among 22 antiviral agents, type I and II recombinant IFNs have the highest potency against SARS-CoV-2 in terms of viral antigen expression, viral load reduction, and plaque reduction ability in VeroE6 cells [60]. These results collectively suggest the potential efficacy of human IFN in suppressing SARS-CoV-2 titers.

4.2. Animal study

Direct IFN treatment or in combination demonstrated protective effects against SARS-CoV or MERS-CoV in experimentally infected mice, marmosets or macaques when administered subcutaneously or intranasally [26,55,61–65]. Haagmans et al. confirmed the *in vivo* anti-SARS-CoV activity of PEG-IFN- α in macaques, wherein PEG-IFN- α effectively hindered viral replication, viral antigen expression by type I pneumocytes, and pulmonary damage [61]. IFN combined with ribavirin and/or lopinavir was effective against MERS-CoV in rhesus macaques [55] and common marmosets [62]. Falzarano et al. demonstrated that IFN- α 2b and ribavirin partly prevented disease progression to pneumonia and reduced viral genome copy number and gene expression levels in rhesus macaques after infection with MERS-CoV [55]. IFN- β 1b and lopinavir, either alone or in combination, could protect marmosets from the adverse effects of MERS-CoV infection in terms of clinical, radiological, and pathological parameters, and the combination showed a better inhibition of viral replication, mitigation of host responses, and improvement of treatment outcomes [62].

However, Sheahan et al. recently reported that IFN- β combined with LPV/RTV exhibited weak antiviral activity against MERS-CoV *in vitro* and *in vivo* when compared with that of remdesivir [48]. The authors also reported that subcutaneous administration of IFN- β every other day failed to decrease the MERS-CoV load and exacerbated the infection in mice [48]. Animal model type, route of administration, and IFN subtype may contribute to inconsistent results among the studies. Therefore, during IFN treatment, the SARS-CoV-2 replication cycle should be evaluated in animal models and/or clinical trials.

5. Clinical data of treatment with IFNs for COVID-19

5.1. IFN treatment of SARS

Patients with SARS are treated with IFNs combined with corticosteroids and antibiotics [8]. In a preliminary open-label, uncontrolled study, in which corticosteroids were used to treat SARS, patients who

also received IFN alfacon-1 had improvement in radiographic lung abnormalities and recovered oxygen saturation levels more quickly than those who received corticosteroids alone [9]. Moreover, the authors reported that patients who took concomitant medications had a lesser increase in creatine kinase levels and a more rapid restoration of lactate dehydrogenase levels to the normal range [9]. However, Zhao et al., in their study, treated patients with SARS with recombinant IFN- α combined with corticosteroids or antibiotics but did not observe the therapeutic benefit of IFN- α [66]. As no new cases have been reported, no further evidence can clarify the therapeutic effects of IFNs on SARS more clearly.

5.2. IFN treatment for MERS

IFNs have been more often used for MERS treatment than for SARS treatment. Routine injections of IFN- β 1b have been recommended as the first choice of treatment [67]. According to the treatment guidelines for MERS stipulated by Korea, PEG-IFN- α 2a can be used at a dosage of 180 μ g/week for 2 weeks unless renal replacement therapy is involved [68].

The combination of IFNs and other antiviral drugs to treat MERS has been described in several case reports [69–71] and retrospective cohort studies [69,72–75]. In a case study, PEG-IFN- α 2a, LPV/RTV, and ribavirin were combined to treat MERS-CoV infection in South Korea, and the results showed satisfactory viral clearance and host survival [70]. Moreover, the survival of patients with severe MERS-CoV infection was improved when treated with IFN- α 2a and ribavirin [74]. The effects of IFNs combined with ribavirin are well documented [57,69,72,76].

Conversely, some studies demonstrated poor or adverse outcomes in patients with MERS who received IFN treatment. For critically ill patients with MERS, different IFN subtypes (rIFN- α 2a, rIFN- α 2b, or rIFN- β 1a) and ribavirin were commonly used, but the drugs did not have any effect on the reduction in 90-day mortality or faster MERS-CoV RNA clearance [77]. Omrani et al. reported that patients with severe MERS-CoV infection who were on mechanical ventilation and who received PEG-IFN- α 2a and ribavirin early in the treatment had a higher 14-day survival rate ($P = .004$), but the difference in the 28-day survival rates between the combination medication and standard care was not statistically significant ($P = .054$) [74]. Another retrospective observational study conducted in Saudi Arabia ($n = 51$) revealed that therapy with IFN- β and mycophenolate mofetil for MERS was associated with a lower mortality rate in univariate analysis but not in multivariate analysis [78]. Although past systematic reviews did not report consensus findings on the efficacy of IFN treatment for MERS [79,80], currently, a randomized controlled trial (RCT) that evaluated 95 people demonstrated that IFN- β combined with LPV/RTV led to a lower 90-day mortality rate than placebo when used within 7 days after symptoms manifestation, which provides strong evidence to support the clinical efficacy of IFN [81].

5.3. IFN treatment for COVID-19

Given the history of IFNs in the treatment of SARS and MERS, IFN- α is used early, especially at treatment initiation, in COVID-19 treatment [82]. In the least official version of the diagnosis and treatment protocol for COVID-19 issued by China, IFN- α is still recommended for antiviral therapy through nebulization (5 million units per time for adults in sterile injection water, twice a day) [83]. This special administration route may have been chosen to avoid the adverse effects of subcutaneous IFN therapies [14], such as adverse reactions due to IFN injections (fever, chills, myalgia and headache), neuropsychiatric problems, and hypersensitivity reactions as observed in a randomized clinical trial [84]. Although accumulating evidence suggests that subcutaneous injection of IFN can increase host survival rate, alleviate symptoms, and decrease hospital length of stay when combined with other medications [84–88], some researchers have thought highly of nebulized IFNs for its higher pulmonary concentration and lower systemic side effects.

Inhalation treatment with IFN- β -1b and LPV/RTV successfully cured patients who had a worsening respiratory status after antibiotic and hydroxychloroquine treatment failed to reverse COVID-19 progression [89]. In a retrospective cohort study of 466 patients, treatment with aerosolized IFN- α 2b within 5 days since hospital admission lowered in-hospital mortality risk [90]. Another double-blind RCT showed that nebulized IFN- β 1b had greater prospects of conferring rapid improvement and recovery [91]. In a systematic review and meta-analysis, several RCTs consistently reported that IFN- β treatment can lower intensive care unit admission risk by 42% and increase the discharge rate by 3.05-fold in patients with COVID-19, thus supporting its use as a safe therapy for COVID-19 [92,93].

However, some trials recommend the avoidance of treatment with IFNs because the IFNs did not demonstrate any synergistic effect when combined with other drugs or therapies [94–96]. Meanwhile, patients with COVID-19 receiving IFN- β -1b treatment may experience elevated anti thyroid peroxidase levels and thyroid dysfunction [97]. Owing to heterogeneous results, treatment with IFNs for COVID-19 warrants further clinical evaluation.

6. IFN efficacy and possible influencing factors

As mentioned earlier, in vitro experiments, animal models, and clinical studies have provided inconsistent evidence on the efficacy of IFNs against MERS, as observed in a systematic review and meta-analysis; the mortality rate of patients who received IFNs (α 2a, α 2b or β 1a) combined with ribavirin was equivalent to that of those who received supportive treatment only [79]. This discrepancy between in vitro and in vivo findings may be attributed to the high EC50/Cmax ratio of the drugs and the late timing of IFN therapy [36,98]. Clinical studies on patients with SARS or MERS report heterogeneous findings, varying in terms of timing of IFN administration, type of patient population assessed, stage of disease, type of IFN used (PEGylated or short-acting), and so on [3,12,36,98]. Similarly, the largest clinical trial of COVID-19 to date, namely, the Solidarity Trial, reported highly controversial results. The trial indicated that IFN- β -1a, remdesivir, hydroxychloroquine, and LPV/RTV regimens were barely conducive to overall mortality or the in-hospital clinical course [95], but as the trial prioritized broad access, people were recruited regardless of significant heterogeneity in therapy implementation, controls (standard care depended on which drugs were used and at what time they were locally available), patient populations, and their clinical conditions. Consequently, whether the results are conclusive remains uncertain [99]. An open-label RCT in which the subjects had more similar background characteristics also supported the finding that IFN- β -1a failed to improve either the clinical status or viral clearance, but the study also confirmed the absence of patients who were in the early phase of COVID-19 [96]. The viral kinetics of respiratory specimens cannot be determined, as most studies conducted on humans are case studies, the tolerability of the IFN regimen by patients with MERS has not been sufficiently evaluated, and the viral load has not been quantified [70]. Furthermore, patients were simultaneously treated with other antivirals, antibiotics, steroids, and invasive or noninvasive ventilation, and IFNs were even administered prophylactically in some cases [69]. We cannot exclude the possibility of spontaneous improvement in these studies, although many patients recovered after receiving IFN antiviral therapy. Thus, the efficacy of IFN treatment from these clinical reports remains controversial, and recommendations for therapy have been barely made.

Several factors described above may affect the clinical outcome of IFN therapy. First, the timing of IFN response induction determines the clinical course of the disease [100]. IFN treatment initiation after 10 days since symptom onset barely exerted any therapeutic effect (Table 1) [68,75], which is probably attributed to the viral load. In the initial stage of infection, the viral load is adequately low for the IFNs to counteract, whereas, in disease exacerbation, the IFN response is thwarted and delayed by the virus, almost always leading to

hyperinflammation and tissue damage [101]. When IFN- β -1b treatment was limited within 7 days, the addition of IFN- β -1b demonstrated safety and superior effects to those of LPV/RTV alone in terms of alleviating symptoms and shortening hospital length of stay in patients with mild-to-moderate COVID-19 [87]. Second, the association between various durations of IFN treatment and clinical outcomes has not yet been validated in double-blind RCTs. Currently, the optimal duration is decided according to the patient's conditions; hence, further well-designed studies evaluating the effects of IFN duration on treatment outcomes in COVID-19 are urgently needed. Third, IFNs are usually been combined with ribavirin or LPV/RTV; thus, the elucidation of data was hindered mainly by various combinations of regimens. Given the lack of convincing clinical evidence, no particular IFN is superior to the other types [72], although a three-arm, open-label RCT reported that IFN- β 1a, rather than IFN- β 1b, significantly shortened the time to clinical improvement [102], and such conclusion should be interpreted with more caution. Another study that suggested the use of IFN- λ for patients with COVID-19 also raised concerns regarding the control of the “cytokine storm” emerging during IFN treatment [103]. In addition, critically ill patients with multiple immunosuppression-associated comorbidities, such as chronic kidney disease, may not benefit from IFN antiviral therapy [72,75,100]. It is therefore necessary to screen all patients and identify who would most benefit from IFN treatment. Further large-scale studies are required, considering the effect of all possible factors including the timing of IFN initiation, IFN treatment duration, IFN type used, and the clinical condition of patients with COVID-19.

7. Conclusion

In this review, we summarized published literature on IFN treatment for COVID-19. As broad-spectrum antiviral agents, IFNs not only directly inhibit viral replication but also activate immune cell subsets, thus providing a potential theoretical benefit for COVID-19 treatment. They are clinically well-established agents and routinely available. Currently available IFN therapies for COVID-19 are mainly adapted from those used for SARS-CoV or MERS-CoV, but in vitro and animal studies have mostly assessed their effects. Regardless of the satisfactory outcomes of these in vitro and nonhuman primate studies, the limited number of case studies or retrospective studies does not provide overall firm evidence concerning the efficacy or safety of IFN therapy per se for CoVs, owing to the lack of well-designed and controlled clinical trials. Moreover, the timing of IFN treatment initiation, the duration of treatment, the IFN type used, and other pivotal factors of IFNs may affect the outcomes of CoV treatment, thereby being worthy of further investigation. In the scenario of the ongoing dangerous CoV outbreak, providing convincing positive or negative findings will be critical for treatment. Thus, the findings of this review may serve as the basis for future clinical trials of IFN treatment for COVID-19.

CRediT authorship contribution statement

Dan Ma: Data curation, Formal analysis, Writing – original draft. **Ximin Wang:** Data curation, Visualization. **Min Li:** Visualization, Investigation, Writing – review & editing. **Chujiao Hu:** Conceptualization, Methodology. **Lei Tang:** Conceptualization, Methodology.

Author contributions

The corresponding authors, LT drew up the outline. DM, ML and XMW searched information to draft the manuscript and redacted it. All authors contributed to the idea for the review and agreed to the submitted version. CJH and LT are the guarantors.

Table 1

Clinical studies of IFN against COVID-19.

Virus infection	Study Design	IFN Treatment Regimen	Combined antiviral drug	Time of IFN initiation	Time of IFN duration	Results	References
SARS-CoV	Retrospective cohort study (n = 22)	IFN alfacon-1 (9 µg/day for the first 2 days and increased to 15 µg/day if no response)	none	A median of 8 days (IQR 4–10) after symptom onset	10 days	Reduced disease-associated impaired oxygen saturation, more rapid resolution of radiographic lung abnormalities and lower levels of creatine kinase	(Loutfy et al., 2003)
SARS-CoV	Cohort study (n = 190)	IFN-α (3 million IU/day)	none	Unknown	Unknown	Less effective than the early use of steroids	(Zhao et al., 2003)
MERS-CoV	Case series (n = 2)	PEG-IFN alfa-2a (180 µg/week)	RBV	The admission day	2 weeks	Recovery	(Khalid et al., 2014)
MERS-CoV	Case report (n = 1)	PEG-IFN-α2a (180 µg/week)	LPV/r and RBV	15 days after cough; 4 days after admission; >4 days after fever	1 week	Recovery	(Kim et al., 2016)
MERS-CoV	Case report (n = 1)	PEG-IFN (180 µg/week)	LPV/r and RBV	13 days after symptom onset	12 days	Died	(Spanakis et al., 2014)
MERS-CoV	Case report (n = 1)	PEG-IFN-α2a (180 µg/week)	RBV	12 days after symptom onset	1 week	Died	(Malik et al., 2016)
MERS-CoV	Retrospective observational study (n = 2)	PEG-IFN-α2a (180 µg/week) or PEG-IFN-α2b (1.5 µg/kg/week)	RBV	11 days after symptom onset	2 or 3 weeks	The 1st patient improved and were discharged while the 2nd patient died	(Ha et al., 2015)
MERS-CoV	Case series (n = 6)	IFN-α2b (180 µg/week)	RBV	1–2 days from onset in survivors and 12–19 days in those who died	2 weeks	3 patients who had comorbid conditions died, and 3 survived	(Khalid et al., 2014)
MERS-CoV	Retrospective observational study (n = 5)	IFN-a2b (100 µg, 130 µg or 144 µg/week)	RBV	A median of 19 days (range 10–22)	100 µg for 2 weeks; 130 µg or 144 µg once	All died	(Al-Tawfiq et al., 2014)
MERS-CoV	Retrospective cohort study (n = 44)	PEG-IFN alfa-2a (180 µg/week)	RBV	A median of 3 days (range 0–8) from diagnosis of MERS-CoV infection.	2 weeks	Improved survival at 14 days but not at 28 days	(Omrani et al., 2014)
MERS-CoV	Retrospective cohort study (n = 32)	IFN-a2a (180 µg/week) Or IFN-b1a (44 µg, 3 times/week)	RBV	Within 1 day following MERS-CoV diagnosis	IFN-a2a for a median duration of 6 days and IFN-b1a for a median duration of 6.5 days	The overall mortality rate was 85% (11/13) in the IFN-a2a group and 64% (7/11) in the IFN-b1a group (P = 0.24); the lack of efficacy of IFN-a2a or IFN-b1a with RBV in treating MERS-CoV	(Shalhoub et al., 2015)
MERS-CoV	Retrospective cohort study (n = 349)	PEG-rIFN-α2b (1.5 µg/kg/week for 2 weeks); PEG-rIFN-α2a (180 µg/week for 2 weeks); rIFN-β1a (44 mg, 3 times/week)	RBV	A median of 2 days from ICU admission, which corresponded to 5 days from hospital admission and 9.0 days from onset of symptoms	A median of 8 days	IFN combined with RBV was not associated with reduction in 90-day mortality or in faster MERS-CoV RNA clearance	(Arabi et al., 2019)
MERS-CoV	Case series (n = 14)	IFN-α2a (180 µg/week)	RBV	6 days from symptom onset	A maximum of 2 weeks	Survival of all patients but no conclusion on IFN efficacy	(Khalid et al., 2016)
MERS-CoV	Observational study (n = 51)	IFN-α or IFN-β (Dosage was unclear)	RBV or mycophenolate mofetil	Unknown	Unknown	IFN-β and mofetil improved survival; The use of IFN-β and IFN-α was associated with survival rates of 78.3%, 75%	(Al et al., 2016)
MERS-CoV	Retrospective cohort (n = 314)	IFN (subtype and dosage were unclear)	RBV	Unknown	Unknown	The use of interferon-RBV was not associated with mortality.	(Alfaraj et al., 2019)
MERS-CoV	Prospective cohort study (n = 8)	IFN-α2b (Dosage was unclear)	RBV and oseltamivir	Unknown	Unknown	5 patients expired, 2 were discharged alive, and 1 remained intubated at the end of the study period	(Al-Hameed et al., 2016)
SARS-CoV-2	Open-label, RCT (n = 81)	IFN-β1a (44 µg/m/dose, 3 times weekly)	HCQ plus LPV/r or atazanavir/ritonavir	A median time of 10 days after symptoms onset (IQR 8–13)	2 weeks	In IFN group, discharge rate on day 14 significantly was rose and 28-day mortality also was reduced	(Davoudi-Monfared et al., 2020)
SARS-CoV-2	non-controlled prospective trial (n=20)	IFN-β1a (44 µg/dose, q.o.d)	HCQ and LPV/r	The admission day (the time before symptom onset < 7 days)	10 days	Virological clearance significantly was decreased and imaging	(Dastan et al., 2020)

(continued on next page)

Table 1 (continued)

Virus infection	Study Design	IFN Treatment Regimen	Combined antiviral drug	Time of IFN initiation	Time of IFN duration	Results	References
SARS-CoV-2	Observational study (n = 41)	IFN- α 2b (3 million IU/time, q.o.d)	LPV/r	For early intervention, IFN was applied within 72 h of admission, while for late intervention, IFN was took after 72 h of admission)	10 days	records was improved significantly in all patients Virus clearance was accelerated and the average time of hospitalization was decreased. Early IFN intervention was better than delayed treatment as indicated by the days of hospitalization	(Wang et al., 2020)
SARS-CoV-2	Open-label, phase 2, RCT (n = 127)	IFN- β 1b (3 doses of 8 million IU, q.o.d)	LPV/r and RBV	A median of 5 days after symptoms onset (IQR 3–7)	2 weeks	No patients died and IFN group was associated with a significant reduction in the duration of viral shedding, symptom alleviation and shorter duration of hospital stay than control group	(Hung et al., 2020)
SARS-CoV-2	Open-label, RCT (n = 66)	IFN- β 1b(250 μ g, q.o.d)	HCQ plus LPV/r or atazanavir/ritonavir	The admission day (a median time of 7 days from symptom onset to admission, IQR 5–9)	2 weeks	With IFN- β 1b treatment, ICU admission rate, the need for invasive mechanical ventilation and 28-day mortality was significant lower	(Rahmani et al., 2020)
SARS-CoV-2	Case series (n = 4)	IFN- β 1b (9.6 million IU/dose/day)	LPV/r or HCQ	The admission day	10–13 days	All were discharged and alive	(Mary et al., 2020)
SARS-CoV-2	Retrospective multicenter cohort study (n = 446)	IFN- α 2b (Dosage was unclear)	LPV/r, or umifenovir	For early IFN, the median time from admission to first dose was 2 days (IQR 1–2), while for late IFN, the median time was 8.5 days (IQR 7–11)	A median of 10 days	As for mortality rate, early administration of IFN exerted better effect than late IFN therapy. And IFN was associated with faster recovery than LPV/r	(Wang et al., 2020)
SARS-CoV-2	Double-blind, phase 2, RCT (n = 98)	IFN- β 1a (6 million IU/dose/day)	Unknown	The admission day (a median duration of 10 days of COVID-19 symptoms, IQR 7–11)	2 weeks	None patients died in IFN group and IFN increased the odds of recovery	(Monk et al., 2021)
SARS-CoV-2	Double-blind, phase 3, RCT(n = 969)	IFN- β 1a (44 μ g/dose, q.o.d)	Unknown	The admission day (a mean duration of 8.7 days from symptoms onset before enrolment)	Up to 4 doses	The addition of IFN- β 1a didn't show more benefit than using remdesivir alone	(Kalil et al., 2021)
SARS-CoV-2	Open-label RCT (n = 11,330)	IFN- β 1a (44 μ g/dose, subcutaneous on the day of randomization and days 3 and 6 or 10 μ g intravenously daily)	With or without lopinavir	The randomization day	6 days	IFN- β 1a didn't benefit COVID-19 patients in terms of overall mortality, initiation of ventilation, and length of hospital stay	(WHO, Solidarity Trial Consortium et al., 2021)
SARS-CoV-2	Open-label RCT (n = 583)	IFN- β 1a (44 μ g/dose)	LPV/r	The randomization day (a median time of 9 days from symptom onset to randomization, IQR 7–12)	On day 1, 3 and 6	No significant effect	(Ader et al., 2021)
SARS-CoV-2	Open-label RCT (n = 60)	IFN- β 1a (12,000 IU) or IFN- β 1b (8 million IU)	HCQ plus LPV/r	The randomization day (a median time of 5 days from symptom onset to randomization, IQR 3–7)	On day 1, 3 and 6	Patients with IFN- β 1a rather than IFN- β 1b showed clinical improvement more quickly than patients in control	(Alavi Darazam et al., 2021)

Abbreviations: LPV/r, lopinavir/ritonavir; RBV, ribavirin; rIFN, recombinant IFN; HCQ, Hydroxychloroquine; IQR, interquartile range.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

References

- [1] P. Weiss, D.R. Murdoch, Clinical course and mortality risk of severe COVID-19, *Lancet* 395 (10229) (2020) 1014–1015, [https://doi.org/10.1016/S0140-6736\(20\)30633-4](https://doi.org/10.1016/S0140-6736(20)30633-4).
- [2] A.M. Zaki, S. van Boheemen, T.M. Bestebroer, A.D. Osterhaus, R.A. Fouchier, Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia, *N. Engl. J. Med.* 367 (19) (2012) 1814–1820, <https://doi.org/10.1056/NEJMoa1211721>.
- [3] D.R. Strayer, R. Dickey, W.A. Carter, Sensitivity of SARS/MERS CoV to interferons and other drugs based on achievable serum concentrations in humans, *Infect. Disord. Drug Targets* 14 (1) (2014) 37–43, <https://doi.org/10.2174/18751526514666140713152858>.

- [4] R. Arbel, Y. Wolff Sagy, M. Hoshen, E. Battat, G. Lavie, R. Sergienko, M. Friger, J. G. Waxman, N. Dagan, R. Balicer, Y. Ben-Shlomo, A. Peretz, S. Yaron, D. Serby, A. Hammerman, D. Netzer, Nirmatrelvir Use and Severe Covid-19 Outcomes during the Omicron Surge, *N. Engl. J. Med.* 387 (9) (2022) 790–798, <https://doi.org/10.1056/NEJMoa2204919>.
- [5] B. Du, H.B. Qiu, X. Zhan, Y.S. Wang, H.Y.J. Kang, X.Y. Li, F. Wang, B. Sun, Z. H. Tong, Pharmacotherapeutics for the New Coronavirus Pneumonia, *Zhonghua Jie He He Hu Xi Za Zhi* 43 (2020) E012, <https://doi.org/10.3760/cma.j.issn.1001-0939.2020.0012>.
- [6] Y. Yang, F. Ye, N. Zhu, W. Wang, Y. Deng, Z. Zhao, W. Tan, Middle East respiratory syndrome coronavirus ORF4b protein inhibits type I interferon production through both cytoplasmic and nuclear targets, *Sci. Rep.* 5 (2015) 17554, <https://doi.org/10.1038/srep17554>.
- [7] G. Li, Y. Fan, Y. Lai, T. Han, Z. Li, P. Zhou, P. Pan, W. Wang, D. Hu, X. Liu, Q. Zhang, J. Wu, Coronavirus infections and immune responses, *J. Med. Virol.* 92 (4) (2020) 424–432, <https://doi.org/10.1002/jmv.25685>.
- [8] S. Shigeta, T. Yamase, Current status of anti-SARS agents, *Antivir. Chem. Chemother.* 16 (1) (2005) 23–31, <https://doi.org/10.1177/095632020501600103>.
- [9] M.R. Loutfy, L.M. Blatt, K.A. Siminovich, S. Ward, B. Wolff, H. Lho, D.H. Pham, H. Deif, E.A. LaMere, M. Chang, K.C. Kain, G.A. Farcas, P. Ferguson, M. Latchford, G. Levy, J.W. Dennis, E.K. Lai, E.N. Fish, Interferon alfacon-1 plus corticosteroids in severe acute respiratory syndrome: a preliminary study, *J. Am. Med. Assoc.* 290 (24) (2003) 3222–3228, <https://doi.org/10.1001/jama.290.24.3222>.
- [10] World Health Organization, 2022. <https://www.emro.who.int/health-topics/mers-cov/mers-outbreaks.html>.
- [11] Johns Hopkins University & Medicine, 2022. <https://coronavirus.jhu.edu/map.html>.
- [12] A.A. Rabaan, S.H. Alahmed, A.M. Bazzi, H.M. Alhani, A review of candidate therapies for Middle East respiratory syndrome from a molecular perspective, *J. Med. Microbiol.* 66 (9) (2017) 1261–1274, <https://doi.org/10.1099/jmm.0.000565>.
- [13] P. Dawson, M.R. Malik, F. Parvez, S.S. Morse, What Have We Learned About Middle East Respiratory Syndrome Coronavirus Emergence in Humans? A Systematic Literature Review, *Vector Borne Zoonotic Dis.* 19 (3) (2019) 174–192, <https://doi.org/10.1089/vbz.2017.2191>.
- [14] G. Li, E. De Clercq, Therapeutic options for the 2019 novel coronavirus (2019-nCoV), *Nat. Rev. Drug Discov.* 19 (3) (2020) 149–150, <https://doi.org/10.1038/d41573-020-00016-0>.
- [15] A. Assiri, A. McGeer, T.M. Perl, C.S. Price, A.A. Al Rabeeah, D.A. Cummings, Z.N. Alabdullatif, M. Assad, A. Almulhim, H. Makhdoom, H. Madani, R. Alhakeem, J. A. Al-Tawfiq, M. Cotten, S.J. Watson, P. Kellam, A.I. Zumla, Z.A. Memish, K.M.-C. I. Team, Hospital outbreak of Middle East respiratory syndrome coronavirus, *N. Engl. J. Med.* 369(5) (2013) 407–16, <https://doi.org/10.1056/NEJMoa1306742>.
- [16] G.A. Almekhlafi, M.M. Albarrak, Y. Mandourah, S. Hassan, A. Alwan, A. Abudayeh, S. Altayyar, M. Mustafa, T. Aldaghestani, A. Alghamdi, A. Talag, M.K. Malik, A.S. Omrani, Y. Sakr, Presentation and outcome of Middle East respiratory syndrome in Saudi intensive care unit patients, *Crit. Care* 20 (1) (2016) 123, <https://doi.org/10.1186/s13054-016-1303-8>.
- [17] L. Kim, S. Garg, A. O'Halloran, M. Whitaker, H. Pham, E.J. Anderson, I. Armistead, N.M. Bennett, L. Billing, K. Como-Sabetti, M. Hill, S. Kim, M. L. Monroe, A. Muse, A.L. Reingold, W. Schaffner, M. Sutton, H.K. Talbot, S. M. Torres, K. Yousey-Hindes, R. Holstein, C. Cummings, L. Brammer, A.J. Hall, A. M. Fry, G.E. Langley, Risk Factors for Intensive Care Unit Admission and In-hospital Mortality Among Hospitalized Adults Identified through the US Coronavirus Disease 2019 (COVID-19)-Associated Hospitalization Surveillance Network (COVID-NET), *Clin. Infect. Dis.* 72 (9) (2021) e206–e214, <https://doi.org/10.1093/cid/ciaa1012>.
- [18] T. Sumi, K. Harada, Immune response to SARS-CoV-2 in severe disease and long COVID-19, *iScience.* 25(8) (2022) 104723, <https://doi.org/10.1016/j.isci.2022.104723>.
- [19] B.X. Wang, E.N. Fish, Global virus outbreaks: Interferons as 1st responders, *Semin. Immunol.* 43 (2019), 101300, <https://doi.org/10.1016/j.smim.2019.101300>.
- [20] J. Zhou, H. Chu, C. Li, B.H. Wong, Z.S. Cheng, V.K. Poon, T. Sun, C.C. Lau, K. K. Wong, J.Y. Chan, J.F. Chan, K.K. To, K.H. Chan, B.J. Zheng, K.Y. Yuen, Active replication of Middle East respiratory syndrome coronavirus and aberrant induction of inflammatory cytokines and chemokines in human macrophages: implications for pathogenesis, *J. Infect Dis* 209 (9) (2014) 1331–1342, <https://doi.org/10.1093/infdis/jit504>.
- [21] B.L. Haagmans, A.D. Osterhaus, Coronaviruses and their therapy, *Antiviral Res.* 71 (2–3) (2006) 397–403, <https://doi.org/10.1016/j.antiviral.2006.05.019>.
- [22] V.A. Scheuplein, J. Seifried, A.H. Malczyk, L. Miller, L. Hocker, J. Vergara-Alert, O. Dolnik, F. Ziebeck, B. Becker, I. Spreitzer, R. Konig, S. Becker, Z. Waibler, M. D. Muhlebach, High secretion of interferons by human plasmacytoid dendritic cells upon recognition of Middle East respiratory syndrome coronavirus, *J. Virol.* 89 (7) (2015) 3859–3869, <https://doi.org/10.1128/JVI.03607-14>.
- [23] Y. Arai, I. Yamanaka, T. Okamoto, A. Isobe, N. Nakai, N. Kamimura, T. Suzuki, T. Daidoji, T. Ono, T. Nakaya, K. Matsumoto, D. Okuzaki, Y. Watanabe, Stimulation of interferon-beta responses by aberrant SARS-CoV-2 small viral RNAs acting as retinoic acid-inducible gene-I agonists, *iScience.* 26(1) (2023) 105742, <https://doi.org/10.1016/j.isci.2022.105742>.
- [24] J. Chen, K. Subbarao, The Immunobiology of SARS*, *Annu. Rev. Immunol.* 25 (2007) 443–472, <https://doi.org/10.1146/annurev.immunol.25.022106.141706>.
- [25] Y. Wang, Y. Gu, D.F. Lewis, X. Gu, K. Brown, C. Lachute, M. Hankins, R.S. Scott, C. Busada, D.B. Cooper, C.E. McCathran, P. Barrilleaux, Cell-type specific distribution and activation of type I IFN pathway molecules at the placental maternal-fetal interface in response to COVID-19 infection, *Front Endocrinol (Lausanne).* 13 (2022), 951388, <https://doi.org/10.3389/fendo.2022.951388>.
- [26] R. Channappanavar, A.R. Fehr, J. Zheng, C. Wohlford-Lenane, J.E. Abrahante, M. Mack, R. Sompallae, P.B. McCray Jr., D.K. Meyerholz, S. Perlman, IFN-I response timing relative to virus replication determines MERS coronavirus infection outcomes, *J. Clin. Invest.* 129 (9) (2019) 3625–3639, <https://doi.org/10.1172/JCI126363>.
- [27] A. Banerjee, N. El-Sayes, P. Budyliowski, R.A. Jacob, D. Richard, H. Maan, J.A. Aguiar, W.L. Demian, K. Baid, M.R. D'Agostino, J.C. Ang, T. Murdza, B.J. Tremblay, S. Afkhami, M. Karimzadeh, A.T. Irving, L. Yip, M. Ostrowski, J.A. Hirota, R. Kozak, T.D. Capellini, M.S. Miller, B. Wang, S. Mubareka, A.J. McGeer, A.G. McArthur, A.C. Doxey, K. Mossman, Experimental and natural evidence of SARS-CoV-2-infection-induced activation of type I interferon responses, *iScience.* 24(5) (2021) 102477, <https://doi.org/10.1016/j.isci.2021.102477>.
- [28] B. Diamond, B.T. Volpe, S. VanPatten, Y. Al Abed, SARS-CoV-2 and interferon blockade, *Mol. Med.* 26(1) (2020) 103, <https://doi.org/10.1186/s10020-020-00231-w>.
- [29] E. Faure, J. Poissy, A. Goffard, C. Fournier, E. Kipnis, M. Titecat, P. Bortolotti, L. Martinez, S. Dubucquoi, R. Desseix, P. Gosset, D. Mathieu, B. Guery, Distinct immune response in two MERS-CoV-infected patients: can we go from bench to bedside? *PLoS One* 9 (2) (2014) e88716, <https://doi.org/10.1371/journal.pone.0088716>.
- [30] F. Ziebeck, M. Weber, M. Eickmann, L. Spiegelberg, A.M. Zaki, M. Matrosovich, S. Becker, F. Weber, Human cell tropism and innate immune system interactions of human respiratory coronavirus EMC compared to those of severe acute respiratory syndrome coronavirus, *J. Virol.* 87 (9) (2013) 5300–5304, <https://doi.org/10.1128/JVI.03496-12>.
- [31] T.R. Tong, SARS coronavirus anti-infectives, *Recent Pat. Anticancer Drug Discov.* 1 (3) (2006) 297–308, <https://doi.org/10.2174/157489106778777637>.
- [32] V. Thiel, F. Weber, Interferon and cytokine responses to SARS-coronavirus infection, *Cytokine Growth Factor Rev.* 19 (2) (2008) 121–132, <https://doi.org/10.1016/j.cytogfr.2008.01.001>.
- [33] X. Lu, J. Pan, J. Tao, D. Guo, SARS-CoV nucleocapsid protein antagonizes IFN-beta response by targeting initial step of IFN-beta induction pathway, and its C-terminal region is critical for the antagonism, *Virus Genes* 42 (1) (2011) 37–45, <https://doi.org/10.1007/s11262-010-0544-x>.
- [34] Y. Yang, L. Zhang, H. Geng, Y. Deng, B. Huang, Y. Guo, Z. Zhao, W. Tan, The structural and accessory proteins M, ORF 4a, ORF 4b, and ORF 5 of Middle East respiratory syndrome coronavirus (MERS-CoV) are potent interferon antagonists, *Protein Cell* 4 (12) (2013) 951–961, <https://doi.org/10.1007/s13238-013-3096-8>.
- [35] V.D. Menachery, A.J. Einfeld, A. Schafer, L. Josset, A.C. Sims, S. Proll, S. Fan, C. Li, G. Neumann, S.C. Tilton, J. Chang, L.E. Gralinski, C. Long, R. Green, C. M. Williams, J. Weiss, M.M. Matzke, B.J. Webb-Robertson, A.A. Schepmoes, A. K. Shukla, T.O. Metz, R.D. Smith, K.M. Waters, M.G. Katze, Y. Kawaoka, R. S. Baric, Pathogenic influenza viruses and coronaviruses utilize similar and contrasting approaches to control interferon-stimulated gene responses, *MBio* 5 (3) (2014) e01174–e01214, <https://doi.org/10.1128/mBio.01174-14>.
- [36] Y. Mo, D. Fisher, A review of treatment modalities for Middle East Respiratory Syndrome, *J. Antimicrob. Chemother.* 71 (12) (2016) 3340–3350, <https://doi.org/10.1093/jac/dkw338>.
- [37] D. Falzarano, E. de Wit, C. Martellaro, J. Callison, V.J. Munster, H. Feldmann, Inhibition of novel beta coronavirus replication by a combination of interferon-alpha2b and ribavirin, *Sci. Rep.* 3 (2013) 1686, <https://doi.org/10.1038/srep01686>.
- [38] H. Zhu, F. Zheng, L. Li, Y. Jin, Y. Luo, Z. Li, J. Zeng, L. Tang, Z. Li, N. Xia, P. Liu, D. Han, Y. Shan, X. Zhu, S. Liu, R. Xie, Y. Chen, W. Liu, L. Liu, X. Xu, J. Wang, H. Yang, X. Shen, X. Jin, F. Cheng, A Chinese host genetic study discovered IFNs and causality of laboratory traits on COVID-19 severity, *iScience.* 24(10) (2021) 103186, <https://doi.org/10.1016/j.isci.2021.103186>.
- [39] E. Kindler, V. Thiel, F. Weber, Interaction of SARS and MERS Coronaviruses with the Antiviral Interferon Response, *Adv. Virus Res.* 96 (2016) 219–243, <https://doi.org/10.1016/bs.aivir.2016.08.006>.
- [40] J. Cinatl, B. Morgenstern, G. Bauer, P. Chandra, H. Rabenau, H.W. Doerr, Treatment of SARS with human interferons, *Lancet* 362 (9380) (2003) 293–294, [https://doi.org/10.1016/s0140-6736\(03\)13973-6](https://doi.org/10.1016/s0140-6736(03)13973-6).
- [41] E.L. Tan, E.E. Ooi, C.Y. Lin, H.C. Tan, A.E. Ling, B. Lim, L.W. Stanton, Inhibition of SARS coronavirus infection in vitro with clinically approved antiviral drugs, *Emerg. Infect. Dis.* 10 (4) (2004) 581–586, <https://doi.org/10.3201/eid1004.030458>.
- [42] L.E. Hensley, L.E. Fritz, P.B. Jahrling, C.L. Karp, J.W. Huggins, T.W. Geisbert, Interferon-beta 1a and SARS coronavirus replication, *Emerg. Infect. Dis.* 10 (2) (2004) 317–319, <https://doi.org/10.3201/eid1002.030482>.
- [43] B. Sainz Jr., E.C. Mossel, C.J. Peters, R.F. Garry, Interferon-beta and interferon-gamma synergistically inhibit the replication of severe acute respiratory syndrome-associated coronavirus (SARS-CoV), *Virology* 329 (1) (2004) 11–17, <https://doi.org/10.1016/j.virol.2004.08.011>.
- [44] B. Morgenstern, M. Michaelis, P.C. Baer, H.W. Doerr, J. Cinatl Jr., Ribavirin and interferon-beta synergistically inhibit SARS-associated coronavirus replication in animal and human cell lines, *Biochem. Biophys. Res. Commun.* 326 (4) (2005) 905–908, <https://doi.org/10.1016/j.bbrc.2004.11.128>.
- [45] U. Stroher, A. DiCaro, Y. Li, J.E. Strong, F. Aoki, F. Plummer, S.M. Jones, H. Feldmann, Severe acute respiratory syndrome-related coronavirus is inhibited by interferon- alpha, *J. Infect Dis* 189 (7) (2004) 1164–1167, <https://doi.org/10.1086/382597>.
- [46] H. Dahl, A. Linde, O. Strannegard, In vitro inhibition of SARS virus replication by human interferons, *Scand. J. Infect. Dis.* 36 (11–12) (2004) 829–831, <https://doi.org/10.1080/00365540410021144>.

- [47] M. Spiegel, A. Pichlmair, E. Muhlberger, O. Haller, F. Weber, The antiviral effect of interferon-beta against SARS-coronavirus is not mediated by MxA protein, *J. Clin. Virol.* 30 (3) (2004) 211–213, <https://doi.org/10.1016/j.jcv.2003.11.013>.
- [48] T.P. Sheahan, A.C. Sims, S.R. Leist, A. Schafer, J. Won, A.J. Brown, S. A. Montgomery, A. Hogg, D. Babusis, M.O. Clarke, J.E. Spahn, L. Bauer, S. Sellers, R. Porter, J.Y. Feng, T. Cihlar, R. Jordan, M.R. Denison, R.S. Baric, Comparative therapeutic efficacy of remdesivir and combination lopinavir, ritonavir, and interferon beta against MERS-CoV, *Nat. Commun.* 11 (1) (2020) 222, <https://doi.org/10.1038/s41467-019-13940-6>.
- [49] M. Bao, Y. Zhang, M. Wan, L. Dai, X. Hu, X. Wu, L. Wang, P. Deng, J. Wang, J. Chen, Y. Liu, Y. Yu, L. Wang, Anti-SARS-CoV immunity induced by a novel CpG oligodeoxynucleotide, *Clin. Immunol.* 118 (2–3) (2006) 180–187, <https://doi.org/10.1016/j.clim.2005.09.014>.
- [50] L. Chen, P. Liu, H. Gao, B. Sun, D. Chao, F. Wang, Y. Zhu, G. Hedenstierna, C. G. Wang, Inhalation of nitric oxide in the treatment of severe acute respiratory syndrome: a rescue trial in Beijing, *Clin. Infect. Dis.* 39 (10) (2004) 1531–1535, <https://doi.org/10.1086/425357>.
- [51] C. Scagnolari, E. Vicenzi, F. Bellomi, M.G. Stillitano, D. Pinna, G. Poli, M. Clementi, F. Dianzani, G. Antonelli, Increased sensitivity of SARS-coronavirus to a combination of human type I and type II interferons, *Antivir. Ther.* 9 (6) (2004) 1003–1011.
- [52] B.J. Hart, J. Dyall, E. Postnikova, H. Zhou, J. Kindrachuk, R.F. Johnson, G. G. Olinger, M.B. Frieman, M.R. Holbrook, P.B. Jahrling, L. Hensley, Interferon-beta and mycophenolic acid are potent inhibitors of Middle East respiratory syndrome coronavirus in cell-based assays, *J. Gen. Virol.* 95 (Pt 3) (2014) 571–577, <https://doi.org/10.1099/vir.0.061911-0>.
- [53] A.H. de Wilde, V.S. Raj, D. Oudshoorn, T.M. Bestebroer, S. van Nieuwkoop, R. Limpens, C.C. Posthuma, Y. van der Meer, M. Barcena, B.L. Haagmans, E. J. Snijder, B.G. van den Hoogen, MERS-coronavirus replication induces severe in vitro cytopathology and is strongly inhibited by cyclosporin A or interferon-alpha treatment, *J. Gen. Virol.* 94 (Pt 8) (2013) 1749–1760, <https://doi.org/10.1099/vir.0.052910-0>.
- [54] Y.J. Jeon, C.H. Gil, A. Jo, J. Won, S. Kim, H.J. Kim, The influence of interferon-lambda on restricting Middle East Respiratory Syndrome Coronavirus replication in the respiratory epithelium, *Antiviral Res.* 180 (2020), 104860, <https://doi.org/10.1016/j.antiviral.2020.104860>.
- [55] D. Falzarano, E. de Wit, A.L. Rasmussen, F. Feldmann, A. Okumura, D.P. Scott, D. Brining, T. Bushmaker, C. Martellaro, L. Baseler, A.G. Benecke, M.G. Katze, V. J. Munster, H. Feldmann, Treatment with interferon-alpha2b and ribavirin improves outcome in MERS-CoV-infected rhesus macaques, *Nat. Med.* 19 (10) (2013) 1313–1317, <https://doi.org/10.1038/nm.3362>.
- [56] H.S. Li, D.I.T. Kuok, M.C. Cheung, M.M.T. Ng, K.C. Ng, K.P.Y. Hui, J.S.M. Peiris, M.C.W. Chan, J.M. Nicholls, Effect of interferon alpha and cyclosporine treatment separately and in combination on Middle East Respiratory Syndrome Coronavirus (MERS-CoV) replication in a human in-vitro and ex-vivo culture model, *Antiviral Res.* 155 (2018) 89–96, <https://doi.org/10.1016/j.antiviral.2018.05.007>.
- [57] J.F. Chan, K.H. Chan, R.Y. Kao, K.K. To, B.J. Zheng, C.P. Li, P.T. Li, J. Dai, F. K. Mok, H. Chen, F.G. Hayden, K.Y. Yuen, Broad-spectrum antivirals for the emerging Middle East respiratory syndrome coronavirus, *J. Infect.* 67 (6) (2013) 606–616, <https://doi.org/10.1016/j.jinf.2013.09.029>.
- [58] E. Mantlo, N. Bukreyeva, J. Maruyama, S. Paessler, C. Huang, Antiviral activities of type I interferons to SARS-CoV-2 infection, *Antiviral Res.* 179 (2020), 104811, <https://doi.org/10.1016/j.antiviral.2020.104811>.
- [59] K.G. Lokugamage, A. Hage, M. de Vries, A.M. Valero-Jimenez, C. Schindewolf, M. Dittmann, R. Rajsbaum, V.D. Menachery, Type I Interferon Susceptibility Distinguishes SARS-CoV-2 from SARS-CoV, *J. Virol.* 94 (23) (2020), <https://doi.org/10.1128/JVI.01410-20>.
- [60] S. Yuan, C.C. Chan, K.K. Chik, J.O. Tsang, R. Liang, J. Cao, K. Tang, J.P. Cai, Z. W. Ye, F. Yin, K.K. To, H. Chu, D.Y. Jin, I.F. Hung, K.Y. Yuen, J.F. Chan, Broad-Spectrum Host-Based Antivirals Targeting the Interferon and Lipogenesis Pathways as Potential Treatment Options for the Pandemic Coronavirus Disease 2019 (COVID-19), *Viruses* 12 (6) (2020), <https://doi.org/10.3390/v12060628>.
- [61] B.L. Haagmans, T. Kuiken, B.E. Martina, R.A. Fouchier, G.F. Rimmelzwaan, G. van Amerongen, D. van Riel, T. de Jong, S. Itamura, K.H. Chan, M. Tashiro, A. D. Osterhaus, Pegylated interferon-alpha protects type 1 pneumocytes against SARS coronavirus infection in macaques, *Nat. Med.* 10 (3) (2004) 290–293, <https://doi.org/10.1038/nm1001>.
- [62] J.F. Chan, Y. Yao, M.L. Yeung, W. Deng, L. Bao, L. Jia, F. Li, C. Xiao, H. Gao, P. Yu, J.P. Cai, H. Chu, J. Zhou, H. Chen, C. Qin, K.Y. Yuen, Treatment With Lopinavir/Ritonavir or Interferon-beta1b Improves Outcome of MERS-CoV Infection in a Nonhuman Primate Model of Common Marmoset, *J Infect Dis* 212 (12) (2015) 1904–1913, <https://doi.org/10.1093/infdis/jiv392>.
- [63] N. De Albuquerque, E. Baig, X. Ma, J. Zhang, W. He, A. Rowe, M. Habal, M. Liu, I. Shalev, G.P. Downey, R. Gorczynski, J. Butany, J. Leibowitz, S.R. Weiss, I. D. McGilvray, M.J. Phillips, E.N. Fish, G.A. Levy, Murine hepatitis virus strain 1 produces a clinically relevant model of severe acute respiratory syndrome in A/J mice, *J. Virol.* 80 (21) (2006) 10382–10394, <https://doi.org/10.1128/JVI.00747-06>.
- [64] J. Zhao, K. Li, C. Wohlford-Lenane, S.S. Agnihothram, C. Flett, J. Zhao, M. J. Gale Jr., R.S. Baric, L. Enjuanes, T. Gallagher, P.B. McCray Jr., S. Perlman, Rapid generation of a mouse model for Middle East respiratory syndrome, *PNAS* 111 (13) (2014) 4970–4975, <https://doi.org/10.1073/pnas.1323279111>.
- [65] R. Channappanavar, L. Lu, S. Xia, L. Du, D.K. Meyerholz, S. Perlman, S. Jiang, Protective Effect of Intranasal Regimens Containing Peptidic Middle East Respiratory Syndrome Coronavirus Fusion Inhibitor Against MERS-CoV Infection, *J Infect Dis* 212 (12) (2015) 1894–1903, <https://doi.org/10.1093/infdis/jiv325>.
- [66] Z. Zhao, F. Zhang, M. Xu, K. Huang, W. Zhong, W. Cai, Z. Yin, S. Huang, Z. Deng, M. Wei, J. Xiong, P.M. Hawkey, Description and clinical treatment of an early outbreak of severe acute respiratory syndrome (SARS) in Guangzhou, PR China, *J Med Microbiol.* 52 (Pt 8) (2003) 715–720, <https://doi.org/10.1099/jmm.0.05320-0>.
- [67] Public Health England, 2017. <https://www.gov.uk/government/publications/mers-cov-clinical-decision-making-support-for-treatment>.
- [68] Y.P. Chong, J.Y. Song, Y.B. Seo, J.P. Choi, H.S. Shin, T. Rapid Response, Antiviral Treatment Guidelines for Middle East Respiratory Syndrome, *Infect Chemother.* 47(3) (2015) 212–22, 10.3947/ic.2015.47.3.212.
- [69] M. Khalid, B. Khan, F. Al Rabiah, R. Alismaili, S. Saleemi, A.M. Rehan-Khalik, I. Weheba, H. Al Abdely, M. Halim, Q.J. Nadri, A.M. Al Dalaan, M. Zeitouni, T. Butt, E. Al Mutairy, Middle Eastern Respiratory Syndrome Corona Virus (MERS CoV): case reports from a tertiary care hospital in Saudi Arabia, *Ann. Saudi Med.* 34 (5) (2014) 396–400, <https://doi.org/10.5144/0256-4947.2014.396>.
- [70] U.J. Kim, E.J. Won, S.J. Kee, S.I. Jung, H.C. Jang, Combination therapy with lopinavir/ritonavir, ribavirin and interferon-alpha for Middle East respiratory syndrome, *Antivir. Ther.* 21 (5) (2016) 455–459, <https://doi.org/10.3851/IMP3002>.
- [71] A. Malik, K.M. El Masry, M. Ravi, F. Sayed, Middle East Respiratory Syndrome Coronavirus during Pregnancy, Abu Dhabi, United Arab Emirates, 2013, *Emerg. Infect. Dis.* 22 (3) (2016) 515–517, <https://doi.org/10.3201/eid2203.151049>.
- [72] S. Shalhoub, F. Farahat, A. Al-Jiffri, R. Simhairi, O. Shamma, N. Siddiqi, A. Mushtaq, IFN-alpha2a or IFN-beta1a in combination with ribavirin to treat Middle East respiratory syndrome coronavirus pneumonia: a retrospective study, *J. Antimicrob. Chemother.* 70 (7) (2015) 2129–2132, <https://doi.org/10.1093/jac/ckv085>.
- [73] I. Khalid, B.M. Alraddadi, Y. Dai, T.J. Khalid, M. Kadri, A.N. Alshukairi, I. A. Qushmaq, Acute Management and Long-Term Survival Among Subjects With Severe Middle East Respiratory Syndrome Coronavirus Pneumonia and ARDS, *Respir. Care* 61 (3) (2016) 340–348, <https://doi.org/10.4187/respcare.04325>.
- [74] A.S. Omrani, M.M. Saad, K. Baig, A. Bahloul, M. Abdul-Matin, A.Y. Alaidaroos, G. A. Almakhlaifi, M.M. Albarak, Z.A. Memish, A.M. Albarak, Ribavirin and interferon alfa-2a for severe Middle East respiratory syndrome coronavirus infection: a retrospective cohort study, *Lancet Infect. Dis.* 14 (11) (2014) 1090–1095, [https://doi.org/10.1016/S1473-3099\(14\)70920-X](https://doi.org/10.1016/S1473-3099(14)70920-X).
- [75] J.A. Al-Tawfiq, H. Momattin, J. Dib, Z.A. Memish, Ribavirin and interferon therapy in patients infected with the Middle East respiratory syndrome coronavirus: an observational study, *Int. J. Infect. Dis.* 20 (2014) 42–46, <https://doi.org/10.1016/j.ijid.2013.12.003>.
- [76] M. Al-Qaseer, The Most Effective Therapeutic Regimen for Patients with Severe Middle East Respiratory Syndrome Coronavirus (MERS-CoV), *Infection, Journal of Infectious Diseases & Therapy.* 03 (04) (2015), <https://doi.org/10.4172/2332-0877.1000223>.
- [77] Y.M. Arabi, S. Shalhoub, Y. Mandourah, F. Al-Hameed, A. Al-Omari, E. Al Qasim, J. Jose, B. Alraddadi, A. Almotairi, K. Al Khatib, A. Abdulmomen, I. Qushmaq, A. A. Sindi, A. Mady, O. Solaiman, R. Al-Raddadi, K. Maghrabi, A. Ragab, G.A. Al Mekhlafi, H.H. Balkhy, A. Al Harthy, A. Kharaba, J.A. Gramish, A.M. Al-Aithan, A. Al-Dawood, L. Merson, F.G. Hayden, R. Fowler, Ribavirin and Interferon Therapy for Critically Ill Patients With Middle East Respiratory Syndrome: A Multicenter Observational Study, *Clin Infect Dis.* 70(9) (2020) 1837–1844, 10.1093/cid/ciz544.
- [78] M. Al Ghamdi, K.M. Alghamdi, Y. Ghandoura, A. Alzahrani, F. Salah, A. Alsulami, M.F. Bawayan, D. Vaidya, T.M. Perl, G. Sood, Treatment outcomes for patients with Middle Eastern Respiratory Syndrome Coronavirus (MERS CoV) infection at a coronavirus referral center in the Kingdom of Saudi Arabia, *BMC Infect Dis.* 16 (2016) 174, 10.1186/s12879-016-1492-4.
- [79] M.E. Morra, L. Van Thanh, M.G. Kamel, A.A. Ghazy, A.M.A. Altibi, L.M. Dat, T.N. X. Thy, N.L. Vuong, M.R. Mostafa, S.I. Ahmed, S.S. Elabd, S. Fathima, T. Le Huy Vu, A.S. Omrani, Z.A. Memish, K. Hirayama, N.T. Huy, Clinical outcomes of current medical approaches for Middle East respiratory syndrome: A systematic review and meta-analysis, *Rev. Med. Virol.* 28 (3) (2018) e1977, DOI: 10.1002/rmv.1977.
- [80] H. Momattin, A.Y. Al-Ali, J.A. Al-Tawfiq, A Systematic Review of therapeutic agents for the treatment of the Middle East Respiratory Syndrome Coronavirus (MERS-CoV), *Travel Med. Infect. Dis.* 30 (2019) 9–18, <https://doi.org/10.1016/j.tmaid.2019.06.012>.
- [81] Y.M. Arabi, A.Y. Asiri, A.M. Assiri, H.H. Balkhy, A. Al Bshabshe, M. Al Jeraisy, Y. Mandourah, M.H.A. Azzam, A.M. Bin Eshaq, S. Al Johani, S. Al Harbi, H.A.A. Jokhdar, A.M. Deeb, Z.A. Memish, J. Jose, S. Ghazal, S. Al Faraj, G.A. Al Mekhlafi, N.M. Sherbeeni, F.E. Elzein, F. Al-Hameed, A. Al Saedi, N.K. Alharbi, R.A. Fowler, F.G. Hayden, A. Al-Dawood, M. Abdelzaher, W. Bajhmom, B.M. AlMutairi, M.A. Hussein, A. Althman, G. Saudi Critical Care Trials, Interferon Beta-1b and Lopinavir-Ritonavir for Middle East Respiratory Syndrome, *N Engl J Med.* 383 (17) (2020) 1645–1656, 10.1056/NEJMoa2015294.
- [82] Y.H. Jin, L. Cai, Z.S. Cheng, H. Cheng, T. Deng, Y.P. Fan, C. Fang, D. Huang, L.Q. Huang, Q. Huang, Y. Han, B. Hu, F. Hu, B.H. Li, Y.R. Li, K. Liang, L.K. Lin, L.S. Luo, J. Ma, L.L. Ma, Z.Y. Peng, Y.B. Pan, Z.Y. Pan, X.Q. Ren, H.M. Sun, Y. Wang, Y.Y. Wang, H. Weng, C.J. Wei, D.F. Wu, J. Xia, Y. Xiong, H.B. Xu, X.M. Yao, Y.F. Yuan, T.S. Ye, X.C. Zhang, Y.W. Zhang, Y.G. Zhang, H.M. Zhang, Y. Zhao, M.J. Zhao, H. Zi, X.T. Zeng, Y.Y. Wang, X.H. Wang, f.t.Z.H.o.W.U.N.C. Management, E-B.M.C.o.C.I.E. Research Team, M. Promotive Association for, C. Health, A rapid advice guideline for the diagnosis and treatment of 2019 novel coronavirus

- (2019-nCoV) infected pneumonia (standard version), *Mil Med Res.* 7(1) (2020) 4, 10.1186/s40779-020-0233-6.
- [83] T.G.O.o.N.H. CommissionOffice, N.T. Administration, Diagnosis and Treatment Protocol for COVID-19 Patients (Tentative 8th Edition), 感染性疾病与免疫(英文). 2(3) (2022) 10.
- [84] E. Davoudi-Monfared, H. Rahmani, H. Khalili, M. Hajiabdolbaghi, M. Salehi, L. Abbasian, H. Kazemzadeh, M.S. Yekaninejad, A Randomized Clinical Trial of the Efficacy and Safety of Interferon beta-1a in Treatment of Severe COVID-19, *Antimicrob. Agents Chemother.* 64 (9) (2020), <https://doi.org/10.1128/AAC.01061-20>.
- [85] F. Dastan, S.A. Nadji, A. Saffaei, M. Marjani, A. Moniri, H. Jamaati, S. M. Hashemian, P. Baghaei, A. Abedini, M. Varahram, S. Yousefian, P. Tabarsi, Subcutaneous administration of interferon beta-1a for COVID-19: A non-controlled prospective trial, *Int. Immunopharmacol.* 85 (2020), 106688, <https://doi.org/10.1016/j.intimp.2020.106688>.
- [86] B. Wang, D. Li, T. Liu, H. Wang, F. Luo, Y. Liu, Subcutaneous injection of IFN alpha-2b for COVID-19: an observational study, *BMC Infect. Dis.* 20 (1) (2020) 723, <https://doi.org/10.1186/s12879-020-05425-5>.
- [87] I.F. Hung, K.C. Lung, E.Y. Tso, R. Liu, T.W. Chung, M.Y. Chu, Y.Y. Ng, J. Lo, J. Chan, A.R. Tam, H.P. Shum, V. Chan, A.K. Wu, K.M. Sin, W.S. Leung, W.L. Law, D.C. Lung, S. Sin, P. Yeung, C.C. Yip, R.R. Zhang, A.Y. Fung, E.Y. Yan, K.H. Leung, J.D. Ip, A.W. Chu, W.M. Chan, A.C. Ng, R. Lee, K. Fung, A. Yeung, T.C. Wu, J. W. Chan, W.W. Yan, W.M. Chan, J.F. Chan, A.K. Lie, O.T. Tsang, V.C. Cheng, T. L. Que, C.S. Lau, K.H. Chan, K.K. To, K.Y. Yuen, Triple combination of interferon beta-1b, lopinavir-ritonavir, and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, randomised, phase 2 trial, *Lancet* 395 (10238) (2020) 1695–1704, [https://doi.org/10.1016/S0140-6736\(20\)31042-4](https://doi.org/10.1016/S0140-6736(20)31042-4).
- [88] H. Rahmani, E. Davoudi-Monfared, A. Nourian, H. Khalili, N. Hajizadeh, N. Z. Jalalabadi, M.R. Fazeli, M. Ghazaeian, M.S. Yekaninejad, Interferon beta-1b in treatment of severe COVID-19: A randomized clinical trial, *Int. Immunopharmacol.* 88 (2020), 106903, <https://doi.org/10.1016/j.intimp.2020.106903>.
- [89] A. Mary, L. Henaut, P.Y. Macq, L. Badoux, A. Cappe, T. Poree, M. Eckes, H. Dupont, M. Brazier, Rationale for COVID-19 Treatment by Nebulized Interferon-beta-1b-Literature Review and Personal Preliminary Experience, *Front. Pharmacol.* 11 (2020), 592543, <https://doi.org/10.3389/fphar.2020.592543>.
- [90] N. Wang, Y. Zhan, L. Zhu, Z. Hou, F. Liu, P. Song, F. Qiu, X. Wang, X. Zou, D. Wan, X. Qian, S. Wang, Y. Guo, H. Yu, M. Cui, G. Tong, Y. Xu, Z. Zheng, Y. Lu, P. Hong, Retrospective Multicenter Cohort Study Shows Early Interferon Therapy Is Associated with Favorable Clinical Responses in COVID-19 Patients, *Cell Host Microbe.* 28(3) (2020) 455–464 e2, 10.1016/j.chom.2020.07.005.
- [91] P.D. Monk, R.J. Marsden, V.J. Tear, J. Brookes, T.N. Batten, M. Mankowski, F.J. Gabbay, D.E. Davies, S.T. Holgate, L.P. Ho, T. Clark, R. Djukanovic, T.M.A. Wilkinson, C.-S.G. Inhaled Interferon Beta, Safety and efficacy of inhaled nebulised interferon beta-1a (SNG001) for treatment of SARS-CoV-2 infection: a randomised, double-blind, placebo-controlled, phase 2 trial, *Lancet Respir Med.* 9 (2) (2021) 196–206, 10.1016/S2213-2600(20)30511-7.
- [92] W.C. Chen, C.K. Hsu, C.Y. Chen, C.C. Lai, S.H. Hung, W.T. Lin, Clinical efficacy and safety of interferon-beta-containing regimens in the treatment of patients with COVID-19: a systematic review and meta-analysis of randomized controlled trials, *Expert Rev. Anti Infect. Ther.* 20 (5) (2022) 741–747, <https://doi.org/10.1080/14787210.2022.2004118>.
- [93] A. Nakhilband, A. Fakhari, H. Azizi, Interferon-beta offers promising avenues to COVID-19 treatment: a systematic review and meta-analysis of clinical trial studies, *Naunyn Schmiedeberg Arch. Pharmacol.* 394 (5) (2021) 829–838, <https://doi.org/10.1007/s00210-021-02061-x>.
- [94] A.C. Kalil, A.K. Mehta, T.F. Patterson, N. Erdmann, C.A. Gomez, M.K. Jain, C.R. Wolfe, G.M. Ruiz-Palacios, S. Kline, J. Regalado Pineda, A.F. Luetkemeyer, M.S. Harkins, P.E.H. Jackson, N.M. Iovine, V.F. Tapson, M.D. Oh, J.A. Whitaker, R.A. Mularski, C.I. Paules, D. Ince, J. Takasaki, D.A. Sweeney, U. Sandkovsky, D.L. Wyles, E. Hohmann, K.A. Grimes, R. Grossberg, M. Laguio-Vila, A.A. Lambert, D. Lopez de Castilla, E. Kim, L. Larson, C.R. Wan, J.J. Traenkner, P.O. Ponce, J.E. Patterson, P.A. Goepfert, T.A. Sofarelli, S. Mocherla, E.R. Ko, A. Ponce de Leon, S. B. Doernberg, R.L. Atmar, R.C. Maves, F. Dangond, J. Ferreira, M. Green, M. Makowski, T. Bonnett, T. Beresnev, V. Ghazaryan, W. Dempsey, S.U. Nayak, L. Dodd, K.M. Tomashek, J.H. Beigel, A.-s.g. members, Efficacy of interferon beta-1a plus remdesivir compared with remdesivir alone in hospitalised adults with COVID-19: a double-blind, randomised, placebo-controlled, phase 3 trial, *Lancet Respir Med.* 9(12) (2021) 1365–1376, 10.1016/S2213-2600(21)00384-2.
- [95] W.H.O.S.T. Consortium, H. Pan, R. Peto, A.M. Henao-Restrepo, M.P. Preziosi, V. Sathiyamoorthy, Q. Abdool Karim, M.M. Alejandria, C. Hernandez Garcia, M.P. Kieny, R. Malekzadeh, S. Murthy, K.S. Reddy, M. Roses Periago, P. Abi Hanna, F. Ader, A.M. Al-Bader, A. Alhasawi, E. Allum, A. Alotaibi, C.A. Alvarez-Moreno, S. Appadoo, A. Asiri, P. Aukrust, A. Barratt-Due, S. Bellani, M. Branca, H.B.C. Cappel-Porter, N. Cerrato, T.S. Chow, N. Como, J. Eustace, P.J. Garcia, S. Godbole, E. Gotuzzo, L. Griskevicius, R. Hamra, M. Hassan, M. Hassany, D. Hutton, I. Irmansyah, L. Jancoriente, J. Kirwan, S. Kumar, P. Lennon, G. Lopardo, P. Lydon, N. Magrini, T. Maguire, S. Manevska, O. Manuel, S. McGinty, M.T. Medina, M.L. Mesa Rubio, M.C. Miranda-Montoya, J. Nel, E.P. Nunes, M. Perola, A. Portoles, M.R. Rasmin, A. Raza, H. Rees, P.P.S. Reges, C.A. Rogers, K. Salami, M.I. Salvadori, N. Sinani, J.A.C. Sterne, M. Stevanovikj, E. Tacconelli, K.A.O. Tikkinen, S. Trelle, H. Zaid, J.A. Rottingen, S. Swaminathan, Repurposed Antiviral Drugs for Covid-19 - Interim WHO Solidarity Trial Results, *N Engl J Med.* 384(6) (2021) 497–511, 10.1056/NEJMoa2023184.
- [96] F. Ader, N. Peiffer-Smadja, J. Poissy, M. Bouscambert-Duchamp, D. Belhadi, A. Diallo, C. Delmas, J. Saillard, A. Dechanet, N. Mercier, A. Dupont, T. Alfaïate, F.X. Lescure, F. Raffi, F. Goehring, A. Kimmoun, S. Jaureguiberry, J. Reigner, S. Nseir, F. Danion, R. Clere-Jehl, K. Bouiller, J.C. Navellou, V. Tolsma, A. Cabie, C. Dubost, J. Courjon, S. Leroy, J. Mootien, R. Gaci, B. Mourvillier, E. Faure, V. Pourcher, S. Gallien, O. Launay, K. Lacombe, J.P. Lanoix, A. Makinson, G. Martin-Blondel, L. Bouadma, E. Botelho-Nevers, A. Gagneux-Brunon, O. Epaulard, L. Piroth, F. Wallet, J.C. Richard, J. Reuter, T. Staub, B. Lina, M. Noret, C. Andrejak, M.P. Le, G. Peytavin, M. Hites, D. Costagliola, Y. Yazdanpanah, C. Burdet, F. Mentre, g. DisCoVeRy study, An open-label randomized controlled trial of the effect of lopinavir/ritonavir, lopinavir/ritonavir plus IFN-beta-1a and hydroxychloroquine in hospitalized patients with COVID-19, *Clin Microbiol Infect.* 27(12) (2021) 1826–1837, 10.1016/j.cmi.2021.05.020.
- [97] D.T.W. Lui, I.F.N. Hung, C.H. Lee, A.C.H. Lee, A.R. Tam, P. Pang, T.Y. Ho, C.Y. Y. Cheung, C.H.Y. Fong, C.Y. Law, K.K.W. To, C.W. Lam, W.S. Chow, Y.C. Woo, K. S.L. Lam, K.C.B. Tan, The Impact of Interferon Beta-1b Therapy on Thyroid Function and Autoimmunity Among COVID-19 Survivors, *Front Endocrinol (Lausanne)*. 12 (2021), 746602, <https://doi.org/10.3389/fendo.2021.746602>.
- [98] A. Zumla, J.F. Chan, E.I. Azhar, D.S. Hui, K.Y. Yuen, Coronaviruses - drug discovery and therapeutic options, *Nat. Rev. Drug Discov.* 15 (5) (2016) 327–347, <https://doi.org/10.1038/nrd.2015.37>.
- [99] D.P. Harrington, L.R. Baden, J.W. Hogan, A. Large, Simple Trial Leading to Complex Questions, *N. Engl. J. Med.* 384 (6) (2021) 576–577, <https://doi.org/10.1056/NEJMe2034294>.
- [100] D. Bencze, T. Fekete, K. Pazmandi, Correlation between Type I Interferon Associated Factors and COVID-19 Severity, *Int. J. Mol. Sci.* 23 (18) (2022), <https://doi.org/10.3390/ijms231810968>.
- [101] A. Park, A. Iwasaki, Type I and Type III Interferons - Induction, Signaling, Evasion, and Application to Combat COVID-19, *Cell Host Microbe* 27 (6) (2020) 870–878, <https://doi.org/10.1016/j.chom.2020.05.008>.
- [102] I. Alavi Darazam, S. Shokouhi, M.A. Pourhoseingholi, S.S. Naghibi Irvani, M. Mokhtari, M. Shabani, M. Amirhosseini, P. Torabianpour, M. Golmohammadi, S. Hashemi, A. Azimi, M.H. Jafarazadeh Maivan, O. Rezaei, A. Zali, M. Hajiesmaeili, H. Shabanpour Dehbsneh, A. Hoseyni Kousha, M. Taleb Shoushtari, N. Khalili, A. Soleymannia, L. Gachkar, A. Khoshkar, Role of interferon therapy in severe COVID-19: the COVIFERON randomized controlled trial, *Sci. Rep.* 11 (1) (2021) 8059, <https://doi.org/10.1038/s41598-021-86859-y>.
- [103] E. Andreaskos, S. Tsiodras, COVID-19: lambda interferon against viral load and hyperinflammation, *EMBO Mol Med.* 12(6) (2020) e12465, 10.15252/emmm.202012465.